

Submitted to AJ

Erupting Cataclysmic Variable Stars in the Nearest Globular Cluster, NGC 6397: Intermediate Polars?¹

Michael M. Shara

Department of Astrophysics, American Museum of Natural History, 79th St. and Central Park West, New York, NY, 10024

mshara@amnh.org

Sasha Hinkley

Department of Astrophysics, American Museum of Natural History, 79th St. and Central Park West, New York, NY, 10024

shinkley@amnh.org

David R. Zurek

Department of Astrophysics, American Museum of Natural History, 79th St. and Central Park West, New York, NY, 10024

dzurek@amnh.org

Christian Knigge

School of Physics and Astronomy, University of Southampton, Southampton SO17 1BJ, United Kingdom

christian@astro.soton.ac.uk

Andrea Dieball

School of Physics and Astronomy, University of Southampton, Southampton SO17 1BJ, United Kingdom

¹Based on observations with the NASA/ESA *Hubble Space Telescope*, obtained from the Data Archive at the Space Telescope Science Institute, which is operated by the Association of Universities for Research in Astronomy, Inc., under NASA contract NAS5-26555. These observations are associated with program #8630.

andrea@astro.soton.ac.uk

ABSTRACT

NGC 6397 is the closest globular cluster, and hence the ideal place to search for faint stellar populations such as cataclysmic variables (CVs). HST and Chandra observers have identified nine certain and likely CVs in this nearby cluster, including several magnetic CV candidates. We have combined our recent UV imagery with archival HST images of NGC 6397 to search for new CV candidates and especially to look for dwarf nova-like eruptive events. We find remarkable and somewhat unexpected dwarf nova-like eruptions of the two well-known cataclysmic systems CV2 and CV3. These two objects have been claimed to be *magnetic* CVs, as indicated by their helium emission-line spectra. Magnetic fields in CVs are usually expected to prevent the disk instability that leads to dwarf nova eruptions. In fact, most field magnetic CVs are observed to *not* undergo eruptions. Our observations of the dwarf nova eruptions of CV2 and CV3 can be reconciled with these objects’ HeII emission lines if both objects are infrequently-erupting intermediate polars, similar to EX Hya. If this is the case for most globular cluster CVs then we can reconcile the many X-ray and UV bright CV candidates seen by Chandra and HST with the very small numbers of erupting dwarf novae observed in cluster cores.

Subject headings: Stars: Cataclysmic Variables— dwarf novae

1. Introduction

Galactic globular clusters are surprisingly rich in luminous X-ray sources (Clark 1975; Katz 1975). Scenarios involving tidal capture by two stars (Fabian, Pringle, & Rees 1975) and/or exchange reactions involving three stars (Hut & Verbunt 1983) are widely believed to be the sources of these strongly interacting binaries in clusters. The remarkable correlation between stellar encounter rate and number of X-ray sources in globular cluster cores (Pooley et al. 2003) supports this viewpoint.

Accreting white dwarf-main sequence star binaries—the cataclysmic variable (CV) stars—are almost certainly produced by the same mechanisms in clusters, and should also be plentiful in globulars (Di Stefano & Rappaport 1994). Perhaps the best cluster to search for this

predicted population is NGC 6397. At a mere 3.3 kpc (Gratton 2003) it is the closest globular cluster to the Sun. Thus even CVs resembling the least luminous known such objects (with $M \sim +11$) in this cluster should be detected with deep HST imagery.

The initial discoveries of multiple X-ray sources (Cool et al. 1993) and $H\alpha$ -bright stars (Cool et al. 1995) in NGC 6397 were rapidly followed by spectrographic identification of essentially certain CVs with hydrogen and helium emission lines (Grindlay et al. 1995; Edmonds et al. 1999). Remarkably, all four CVs in this cluster with HST spectra show significant, and in three of four cases, prominent HeII 4686. This line is seen almost exclusively in magnetic CVs and nova-like variables (Williams 1983; Echevarria 1988), prompting Grindlay et al. (1995) and Grindlay (1999) to suggest that magnetic CVs might well be the dominant CV population in globular clusters.

About half of all known field CVs (Downes et al. 2001) are dwarf novae (DN). Most field DN have been discovered because they undergo 2-5 magnitude outbursts every few weeks to months (Warner 1995). In contrast, strongly magnetic CVs are *not* observed to undergo DN outbursts. This is because DN eruptions are likely caused by a disk instability (Mineshige & Osaki 1983), and the accretion disks of magnetic CVs have been truncated or are absent.

Since large archival HST datasets of globular cluster images are available, we have been systematically looking for erupting dwarf novae in the cores of all such clusters (Shara et al. 1996). Here we report the results of a search of NGC 6397 to determine if any of the magnetic CV candidates undergo eruptions. Remarkably we do find dwarf nova eruptions of two of the putative NGC 6397 magnetic CVs.

2. Observations

The Hubble Space Telescope has imaged NGC 6397 during 5 separate epochs from 1996 to 2003. Our own 2003 datasets focused on STIS UV imagery, heretofore unavailable and particularly useful in detecting CVs. The dates of observation, PI and number of HST program, filters used, number of frames and total exposure time in each filter are given in the observing log which is Table 1.

3. Multi-Wavelength Images and Light Curves of Two Dwarf Novae

We have carried out visual inspection and aperture photometry of each of the nine known CVs (referred to by Grindlay et al. (2001) as CV1 through CV9) in every available archival HST image. While a modest level of variability (up to a few tenths of a magnitude)

is detectable in all nine CVs, and two of the objects (CV1 and CV6) now have measured photometric periods (Kaluzny & Thompson 2003), none is yet reported to show a dwarf nova-like outburst.

The HST images of CV2 and CV3 are shown as photo-montages in Figures 1 and 2, respectively. The median brightness in every epoch, and in every available filter is shown with mean filter wavelength running from blue (left) to red (right). We also show the 5-epoch light-curves of these two objects in Figure 3.

Each CV has been imaged in the WFPC2 F814W filter in epochs 1, 2 and 3, and in the STIS CCDclear filter in epochs 4 and 5. We have normalized epochs 1-3 with the first observation of each CV in epoch 1, and epochs 4-5 with the first observation in epoch 4.

The key result of this paper is that the putative magnetic cataclysmic variables CV2 and CV3 are clearly detected as erupting dwarf novae.

CV2 is seen to brighten (in epoch 5) in the Space Telescope Imaging Spectrograph (STIS) UV filter images. It is 2.7 magnitudes brighter than it was in epoch 4.

CV3 is seen in eruption in the second epoch in visible and near infrared pass-band images taken with the HST Wide Field and Planetary Camera (WFPC2). It is 1.8 magnitudes brighter in eruption than in quiescence. It is also seen in an intermediate brightness state (between eruption and quiescence) in blue and near-UV WFPC2 images in epoch 3. The rise time of CV3 in epoch 2 (about 2 days) is well in accord with field dwarf nova rise times.

4. Are the Erupting Systems Magnetic?

The HST spectra of CV1, CV2 and CV3 are presented in Grindlay et al. (1995); that of CV4 is given in Edmonds et al. (1999). HeII emission with equivalent widths in the range 6-15 Å appear in the spectra of all four objects, as do strong Balmer lines. In particular, Edmonds et al. (1999) found for CV2 and CV3, respectively, HeII 4686 equivalent widths of 11 and 15 Å. The H β equivalent widths of these same two objects are 32 and 59 Å, respectively.

The disks in all four objects are faint ($M \sim 8 - 10$). Edmonds et al. (1999) has carefully compared the disks and HeII line ratios of CVs 1-4 with those of other cataclysmic systems. We refer the interested reader to that careful and exhaustive analysis, simply quoting here Edmond et al.’s conclusions regarding CVs 1-3:

“They do not appear to be recent novae or nova-likes because of their faint disks (with

extra evidence from their HeII 4686 Å line ratios), nor do they appear to be dwarf novae because they have moderately strong HeII lines. The final option is magnetic systems. To conclude, CVs 1-3 do not appear to be dwarf novae, but they could be DQ Her-type systems. A possible alternative to the DQ Her hypothesis is that some of the NGC 6397 CVs are old novae (possibly in deep hibernation between outbursts; see Shara et al. 1986).”

Figures 1 and 2 of this paper convincingly demonstrate that CV2 and CV3 are, in fact, erupting dwarf novae despite their prominent HeII emission.

Intensive monitoring with HST is essential to determine if all of the CV candidates in NGC 6397—and other globular clusters—eventually erupt as dwarf novae.

The hibernation scenario of CVs (Shara et al. 1986) postulates that old novae eventually display dwarf-nova like outbursts. One old classical nova in the globular cluster M80 has been recovered (Shara & Drissen 1995) at dwarf nova-like luminosity, and one recent erupting classical nova in a globular cluster (of the giant elliptical galaxy M87) has been reported (Shara et al. 2004). CV2 and CV3 could be hibernating old novae, but we have, at present, no observational proof that they once underwent nova eruptions.

Edmonds’ suggestion that the NGC 6397 CVs might be DQ Her-like magnetic systems (also known as IPs or Intermediate Polars) might seem to be disproved with the observed dwarf nova eruptions of CV2 and CV3. However, there are a few well-documented cases of IPs that undergo dwarf nova eruptions, including GK Per, EX Hya, XY Ari, DO Dra, TV Col, HT Cam and V1223 Sgr.

Most dwarf novae show some HeII emission (see e.g. Patterson & Raymond (1985) for a compilation), so CV2 and CV3 are certainly not unique in this respect. A comparison of the HeII and H β line strengths for the CVs in NGC 6397 and for field IPs is presented by Grindlay (1999). Both the line strengths and line ratios of HeII and H β for CV2 and CV3 are comparable to those of other low mass transfer rate IPs, with Ex Hya being a particularly good match.

Angelini & Verbunt (1989) have presented a model for dwarf nova-like outbursts in magnetic CVs, which they apply to TV Col and GK Per. If the white dwarf in a cataclysmic variable has a strong magnetic field, the inner region of the accretion disk is disrupted. They found that disruption of the inner disk region leads to shorter outbursts, and, for disk instabilities starting near the inner disk edge, to much longer intervals between outbursts.

The outbursts of most IPs tend to be rare, short and of only moderate amplitude. As an example, Hellier et al. (1989) has noted that the EX Hya outbursts occur at ~ 2 year intervals, last less than 4 days and are of only ~ 3 magnitude amplitude. If the many

globular cluster CV candidates identified by Chandra (Grindlay, Heinke, Edmonds, & Murray 2001) and HST (Knigge et al. 2002) are also intermediate polars then the remarkably small numbers of erupting dwarf novae found by HST in globular clusters (Shara et al 1996) can be explained. In particular, a population of even dozens of IPs will only occasionally show a dwarf nova near maximum light. Under this scenario we can predict that more of these CVs will eventually be seen to erupt if individual clusters are searched on dozens or hundreds of occasions.

5. Conclusions

We have observed NGC 6397 with HST during 5 separate epochs. Two of the four spectrographically confirmed CVs are observed to undergo dwarf nova eruptions with amplitudes of at least 1.8 and 2.7 magnitudes. These eruptions were somewhat unexpected, as both objects display moderately strong HeII 4686 emission lines, generally associated with non-erupting, magnetic CVs. The dwarf nova-like eruptions reported in this paper can occur in intermediate polars, suggesting that some of the NGC 6397 CVs (and by implication some of the CVs in other globular clusters) are indeed magnetic systems. The apparent rarity of erupting dwarf novae in globular clusters with large CV candidate populations is reconciled if those CVs are infrequently erupting intermediate polars.

Support for program #8630 was provided by NASA through a grant from the Space Telescope Science Institute, which is operated by the Association of Universities for Research in Astronomy, Inc., under NASA contract NAS 5-26555.”

REFERENCES

- Angelini, L. & Verbunt, F. 1989 MNRAS, 238, 697
- Clark, G. W. 1975, ApJ, 199, L143
- Cool, A.M., Grindlay, J.E., Krockenberger, M., & Bailyn, C.D. 1993 ApJ, 410, L103
- Cool, A.M., Grindlay, J.E., Cohn, H.N., Lugger, P.M., & Slavin, S.D. 1995 ApJ, 439, 695
- Di Stefano, R. & Rappaport, S. 1994, ApJ, 423, 274
- Downes, R. A., Webbink, R. F., Shara, M. M., Ritter, H., Kolb, U., & Duerbeck, H. W. 2001, PASP, 113, 764

- Echevarria, J. ApJ, 233, 513
- Edmonds, P.D., Grindlay, J.E., Cool, A. M., Cohn, H., Lugger, P., & Bailyn, C. 1999 ApJ, 516, 250
- Fabian, A. C., Pringle, J. E., & Rees, M. J. 1975, MNRAS, 172, 15P
- Gratton, R.G., Bragaglia, A., Carretta, E., Clementini, G., Desidera, S., Grundahl, F. & Lucatello, S. 2003 A&A, 408, 529
- Grindlay, J. E. 1999, Astronomical Society of the Pacific Conference Series, 157, 377
- Grindlay, J. E., Cool, A. M., Callanan, P. J., Bailyn, C. D., Cohn, H. N., & Lugger, P. M. 1995, ApJ, 455, L47
- Grindlay, J. E., Heinke, C., Edmonds, P. D., & Murray, S. S. 2001, Science, 292, 2290
- Grindlay, J. E., Heinke, C. O., Edmonds, P. D., Murray, S. S., & Cool, A. M. 2001, ApJ, 563, L53
- Hellier, C., Mason, K., Smale, A., Corbet, R., O'Donoghue, D., Barrett, P., & Warner, B. 1989 MNRAS, 238, 1107
- Hut, P., & Verbunt, F. 1983, Nature, 301, 587
- Kaluzny, J. & Thompson, I. 2003 AJ, 125, 2534
- Katz, J. I. 1975, Nature, 253, 698
- Mineshige, S. & Osaki, Y. 1983, PASP, 35, 377
- Knigge, C., Zurek, D., Shara, M. & Long, K. 2002 ApJ, 579, 752
- Patterson, J. & Raymond, J. C. 1985 ApJ, 292, 535
- Pooley, D., et al. 2003, ApJ, 591, L131
- Shara, M., Bergeron, L., Gilliland, R., Saha, A., & Petro, L. 1996 ApJ, 471, 804
- Shara, M., Livio, M., Moffat, A.F.J. & Orio, M. 1986 ApJ, 311, 163
- Shara, M. M. & Drissen, L. 1995, ApJ, 448, 203
- Shara, M. M., Zurek, D. R., Baltz, E. A., Lauer, T. R., & Silk, J. 2004, ApJ, 605, L117

Warner, B. 1995, Cambridge Astrophysics Series, Cambridge, New York: Cambridge University Press, —c1995,

Williams, G. 1983 ApJS, 53, 523




Fig. 1.— Matrix of HST images of the field of CV2 in NGC6397. North and East are as indicated, and each postage-stamp is $1.55''$ on a side. The top labels indicate the HST filter name, and are organized roughly in increasing wavelength from left (blue) to right (red). CV2 is indicated in each image.




Fig. 2.— Same as Fig. 1, but for CV3.




Fig. 3.— A five-epoch lightcurve for CV2 (top) and CV3 (bottom). The first three epochs show WFPC2 F814W photometry while the last two show the STIS clear CCD data. The magnitudes are purely differential (see text), and the MJDs are indicated for each epoch.

Table 1. Table of NGC6397 observations: Listed are the number of useable observations at each date, as well as their exposure time.

WFPC2 Epochs	PI/Prog. ID	F336W	F439W	F555W	F656N	F675W	F814W
1: 3/6/96	King #5929	700s,6×500s, 28×400s,2×80s, 2×10s	4×500s, 16×400s, 2×80s,2×10s	6×40s,8s,1s			2×40s,8s,1s
2: 4/3/99	Grindlay #7335			24×40s,2×8s 2×1s	1000s,4×900s 18×800s,7×700s 2×140s	38×40s 2×8s,2×1s	24×40s 2×8s,2×1s
3: 11/4/01	Noll #9313	3×160s	2×160s				3×40s
STIS Epochs	PI/Prog. ID	Clear CCD	CCD LP	Clear MAMA	F25QTZ	F25SRF2	
4: 3/28/03	Shara #8630	3×180s	3×180s		533s,865s 6×600s		
5: 5/16/03	“”	3×180s	3×180s	3×1000s		2×600s,883s 2×1021s	